

# Population Health Insights Guide to Help Identify Patients for Shingles Vaccination and Evaluate Shingles Immunization Rates in Your Health System



---

## Indication

SHINGRIX is a vaccine indicated for prevention of herpes zoster (HZ) (shingles):

- in adults aged 50 years and older.
- in adults aged 18 years and older who are or will be at increased risk of HZ due to immunodeficiency or immunosuppression caused by known disease or therapy.

SHINGRIX is not indicated for prevention of primary varicella infection (chickenpox).

## Important Safety Information

- SHINGRIX is contraindicated in anyone with a history of a severe allergic reaction (eg, anaphylaxis) to any component of the vaccine or after a previous dose of SHINGRIX
- Review immunization history for possible vaccine sensitivity and previous vaccination-related adverse reactions. Appropriate medical treatment and supervision must be available to manage possible anaphylactic reactions following administration of SHINGRIX

Please see **Important Safety Information for SHINGRIX** on [page 6](#) and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](http://SHINGRIXHCP.com).

# Table of Contents

<b>01.</b>	Guide Overview .....	3
<b>02.</b>	Background .....	4
<b>03.</b>	Indication and Important Safety Information.....	6
<b>04.</b>	Oracle Health EHR HealthIntent Population Health Suite High Level Overview.....	7
<b>05.</b>	Overview of HealthRegistries Within the HealthIntent Module.....	8
<b>06.</b>	Using HealthRegistry to Track Shingles Vaccinations at the System Level .....	9
<b>07.</b>	Technical Instructions for Configuring HealthRegistries to Help Identify Patients Potentially Eligible for Shingles Immunization per ACIP Recommendations.....	10
<b>08.</b>	Technical Instructions for Modifying Registry Inclusion Rules for Registries .....	11
<b>09.</b>	How to Configure Registry Metrics/Key Performance Indicators to Monitor Population Health Metrics in Oracle Health.....	12
<b>10.</b>	Key Population Health Configuration Use Cases.....	21
<b>11.</b>	Disclaimers .....	24
<b>12.</b>	Notes.....	25

Please see Important Safety Information for SHINGRIX on [page 6](#) and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](#).



# 01. Guide Overview

This Population Health Insights Guide is intended to help health systems make configuration updates in the Oracle Health (formerly Cerner) EHR Population Health Suite (HealthIntent) to generate reports that may identify useful insights to support improving shingles immunization rates. The guide provides suggestions and instructions to monitor and manage shingles immunization rates and associated population health metrics across the health system.

This guide will not work for other conditions, treatments, therapeutic areas, or for other EHR systems.

This guide provides specific considerations for:

- Identifying patients with shingles immunization care gaps
- Using Oracle Health-released registries to provide visibility into immunization care gaps
- Using clinical decision support tools to create care gap awareness in the EHR for clinicians

Additionally, this guide will provide an overview of the HealthIntent module, commonly used features of HealthIntent, and technical instructions to configure these features across the registries, reporting tools, and health maintenance features within the HealthIntent solution suite.

The processes outlined in this piece are variable, and not all steps will apply to every customer. Any steps or settings that are not part of a customer's standard process should be excluded or modified accordingly. Any questions should be directed to the appropriate service provider. The practice is solely responsible for implementing, testing, monitoring, and operating of any EHR tools.

This guide is designed for organizations using the most recent Oracle Health code set. Some configurations may be required by a clinical analyst.

## Key Terms Used in This Guide

- ACIP = Advisory Committee on Immunization Practices
- AIS = Adult Immunization Status
- HCP = healthcare provider
- HZ = herpes zoster
- IC = immunocompromised
- YOA = years of age
- DA2 = Discern Analytics 2

**Please see Important Safety Information for SHINGRIX on page 6 and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](#).**



## 02. Background

### Importance of Managing Adult Immunization to Support Population Health at the Health System Level

The Oracle Health EHR system offers functionality for health systems to manage immunizations at the system or population level by supporting a health system's existing population health strategy. After a health system sets a strategy, which can include improving immunization rates amongst vulnerable populations and standardizing immunization processes, tactics can be set at both the health system and local levels to reach these goals. HealthIntent provides critical tools to support both the strategy and tactics.

Individual tactics—for example, creating a list of patients with immunization care gaps—can help provide valuable care and insights into what is working well at both the health system and individual provider levels. A system-wide approach is suggested to create economies of scale and reduce any outliers that may negatively affect the health system's shingles immunization efforts.

This guide provides instructions on how to help identify patients who may fall within SHINGRIX's indications. These include patients who are  $\geq 50$  YOA and adults aged 18 years and older who are or will be at increased risk of HZ due to immunodeficiency or immunosuppression caused by known disease or therapy.

The following studies supported the approval of the indications described above.

Clinical Trial Experience in Adults 50 Years and Older: The efficacy and safety of SHINGRIX in adults 50 years and older was assessed in two studies. Study 1 (NCT01165177) included adults  $\geq 50$  YOA and Study 2 (NCT01165229) included adults  $\geq 70$  YOA. These studies excluded, among others, subjects who were immunocompromised, had a history of previous HZ, were vaccinated against varicella or HZ, and patients whose survival was not expected to be at least 4 years or with conditions that might interfere with study evaluation.<sup>1</sup>

Clinical Trial Experience in Immunocompromised Adults Aged 18 Years and Older: The safety of SHINGRIX was evaluated in 6 placebo-controlled clinical studies that enrolled subjects aged 18 years and older from 5 different immunodeficient or immunosuppressed (referred to as immunocompromised) populations. These populations included autologous hematopoietic stem cell transplant (aHSCT) recipients (NCT01610414 & NCT00920218), patients with hematologic malignancies who received the first dose of SHINGRIX during or within 6 months of completing immunosuppressive therapy (NCT01767467), renal transplant recipients (NCT02058589), patients with solid malignant tumors receiving chemotherapy (NCT01798056), and patients with HIV (NCT01165203).<sup>1</sup>

Please see Important Safety Information for SHINGRIX on page 6 and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](#).



## 02. Background (continued)

The efficacy of SHINGRIX was evaluated in immunocompromised patients  $\geq 18$  YOA who received an auHSCT (NCT01610414) and was calculated post hoc in another randomized, placebo-controlled, observer-blind study in subjects with hematologic malignancies (NCT01767467).<sup>1</sup>

Dosing information for SHINGRIX<sup>1</sup>

SHINGRIX is administered as a 2-dose series according to the following schedules:

- A first dose at Month 0 followed by a second dose administered 2 to 6 months later.
- For individuals who are or will be immunodeficient or immunosuppressed and who would benefit from a shorter vaccination schedule: A first dose at Month 0 followed by a second dose administered 1 to 2 months later.

Please see Important Safety Information for SHINGRIX on [page 6](#) and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](http://SHINGRIXHCP.com).



## 03. Indication and Important Safety Information

### Indication

SHINGRIX is a vaccine indicated for prevention of herpes zoster (HZ) (shingles):

- in adults aged 50 years and older.
- in adults aged 18 years and older who are or will be at increased risk of HZ due to immunodeficiency or immunosuppression caused by known disease or therapy.

SHINGRIX is not indicated for prevention of primary varicella infection (chickenpox).

### Important Safety Information

- SHINGRIX is contraindicated in anyone with a history of a severe allergic reaction (eg, anaphylaxis) to any component of the vaccine or after a previous dose of SHINGRIX
- Review immunization history for possible vaccine sensitivity and previous vaccination-related adverse reactions. Appropriate medical treatment and supervision must be available to manage possible anaphylactic reactions following administration of SHINGRIX
- In a postmarketing observational study, an increased risk of Guillain-Barré syndrome was observed during the 42 days following vaccination with SHINGRIX
- Syncope (fainting) can be associated with the administration of injectable vaccines, including SHINGRIX. Procedures should be in place to avoid falling injury and to restore cerebral perfusion following syncope
- Solicited local adverse reactions reported in individuals aged 50 years and older were pain (78%), redness (38%), and swelling (26%)
- Solicited general adverse reactions reported in individuals aged 50 years and older were myalgia (45%), fatigue (45%), headache (38%), shivering (27%), fever (21%), and gastrointestinal symptoms (17%)
- Solicited local adverse reactions reported in autologous hematopoietic stem cell transplant recipients (aged 18 to 49 and  $\geq 50$  years of age) were pain (88% and 83%), redness (30% and 35%), and swelling (21% and 18%)
- Solicited general adverse reactions reported in autologous hematopoietic stem cell transplant recipients (aged 18 to 49 and  $\geq 50$  years of age) were fatigue (64% and 54%), myalgia (58% and 52%), headache (44% and 30%), gastrointestinal symptoms (21% and 28%), shivering (31% and 25%), and fever (28% and 18%)
- The data are insufficient to establish if there is vaccine-associated risk with SHINGRIX in pregnant women
- It is not known whether SHINGRIX is excreted in human milk. Data are not available to assess the effects of SHINGRIX on the breastfed infant or on milk production/excretion
- Vaccination with SHINGRIX may not result in protection of all vaccine recipients

Please see full [Prescribing Information](#), available at [SHINGRIXHCP.com](#).



---

## 04. Oracle Health EHR HealthIntent Population Health Suite High Level Overview

---

HealthIntent, Oracle Health's population health management solution, includes:

- Comprehensive care management tools
- Advanced reporting capabilities for actionable insights
- Engagement solutions for both providers and patients
- Tools to empower patient activation
- Support for value-based care initiatives

Health systems can utilize HealthIntent to help improve shingles immunization rates.

Please see Important Safety Information for SHINGRIX on [page 6](#) and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](http://SHINGRIXHCP.com).



## 05. Overview of HealthRegistries Within the HealthIntent Module

A critical component of HealthIntent is **the HealthRegistry solution**. Registries are commonly used to drive clinical outcomes at the health system level, provide data insights, and expose clinical variation to help guide the health system to meet strategic imperatives for achieving shingles vaccination for all appropriate patients.

Registries allow health organizations to identify subsets of their patient population. Registries are organized around a common health element(s) known as **registry inclusion rules**. Common inclusion rules include age, chronic conditions, diagnoses, health measures, immunization status, or other clinical factors. Patients who meet the selected inclusion rule for a particular registry are automatically included in the registry.

Since registries are dynamic, patients who no longer meet the inclusion metric are excluded from the registry, and new patients may be introduced. This is handled by a back-end program and does not require providers to actively enroll patients in a registry.

Oracle Health provides many commonly used health registries to health systems; the most relevant registry for immunizations in the Oracle Health EHR system is the **Wellness Registry**. However, organizations can modify registries and their inclusion rules to meet their desired specific population health programs.

It is important to note that registries can consume data feeds from outside organizations if desired. For immunizations, consider identifying if a state immunization registry is available to feed historic administered immunizations. Adding this data feed can help provide a more complete immunization history to providers, and the immunization rules acted upon by the registry can forecast any immunization(s) due.

Defining registries and registry inclusion factors, components, and outcomes are critical elements of a population health strategy.

**Registry metrics and supporting facts** are components used to collect data that are pertinent to the patients in the defined population registry. Once a metric and supporting fact is associated with a registry, reports can be created from the population subset, and these metrics can be used to show the captured data to gain insight on the patient's condition.

Proposed registry metrics for shingles immunization registries and technical instructions for configuring them can be found in later sections of this guide.

Please see **Important Safety Information for SHINGRIX on page 6** and full **Prescribing Information**, also available at [SHINGRIXHCP.com](https://www.shingrixhcp.com).



## 06. Using HealtheRegistry to Track Shingles Vaccinations at the System Level

Population health insights and registries at the health system level may offer:

- The opportunity to evaluate health system immunization rates
- The ability to help identify and monitor sites within the health system that are not achieving immunization goals
- Shingles-specific clinical decision tools, such as Health Maintenance Alerts and Discern Alerts, creating a standards-based approach to addressing shingles
- The ability to utilize social determinants of health to help identify barriers and help develop impactful shingles immunization efforts

As health systems provide longitudinal healthcare to its members, setting up immunization tactics at the system level can potentially add supplementary touchpoints with patients (expanding from primary care-based encounters to acute and other care settings).

Shingles immunization campaigns can be launched at the local level and expanded and scaled to the health system level if desired. Consider modifying the patient report and Health Maintenance Alert scope from a local instance to the health system when scaling up.

**Please see Important Safety Information for SHINGRIX on [page 6](#) and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](http://SHINGRIXHCP.com).**



---

## 07. Technical Instructions for Configuring HealtheRegistries to Help Identify Patients Potentially Eligible for Shingles Immunization per ACIP Recommendations

---

Registries are based on a general inclusion rule and may contain multiple metrics to define outcomes related to the common registry element. For immunizations, the inclusion rule is age, and multiple wellness age-based registries are available.

The “Comprehensive Adult Wellness” Registry is the default HealtheRegistry that includes all health system patients used for setting adult immunization expectations (additional Registries include the same Measure, for example, the “Senior Wellness” and “Adult Immunization” Registries). Patients who meet ACIP recommendations will automatically be assigned when HealtheRegistries and Health Maintenance are aligned in Oracle Health when they meet the immunization criteria.

Please note that when not electing to use Health Maintenance Alerts, manual Discern Alerts can be created and used. This option would require manual configuration and a maintenance burden on the health system to manage all immunization recommendations and preventive health maintenance recommendations from the US Preventive Services Task Force.

**Please see Important Safety Information for SHINGRIX on [page 6](#) and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](#).**



## 08. Technical Instructions for Modifying Registry Inclusion Rules for Registries

The registry inclusion rule determines who should be part of the registry. This is usually a condition or disease, whereas age is used for the Wellness Registry.

Registry changes can be made in the HealtheRegistries administrator tool if desired. Select the Shingles vaccination Measure Name and select the Inclusion, Exclusion, and Met Criteria as desired.

The integration of the HealtheRegistries and Health Maintenance requires an association in the healthexp tool to appropriately reflect the immunizations in the HealtheRegistries tab of the Recommendations section in PowerChart. To set up the association:

- 1. Select the shingles schedule** in the healthexp tool.
- 2. Check the radio button** next to Associate to a HealtheRegistry measure on the Expectation screen and **select the shingles series**.
- 3. Click Validate** and then **Commit** to save the information.

**NOTE:** The Wellness Registry may not capture all patient vaccinations outside the health system. Therefore, it's important for HCPs to review patient immunization history and update vaccination records as appropriate.

Please see Important Safety Information for SHINGRIX on [page 6](#) and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](http://SHINGRIXHCP.com).



## 09. How to Configure Registry Metrics/Key Performance Indicators to Monitor Population Health Metrics in Oracle Health

Registry metrics provide insight into what needs to be measured for the subset of patients in the registry.

Oracle Health-released registries come with multiple registry metrics. Metric rules can be edited, modified, or added.

To create or modify the configuration logic, logic expression and Enabled/Disabled status as desired.

For the Immunization registry metrics, consider the following Key Performance Indicators (KPIs) and related parameters. These KPIs serve as examples of reports a health system may find useful for tracking shingles immunizations. However, each health system can customize and configure KPIs based on its specific population health priorities and initiatives.

REGISTRY METRIC	KPI	RULE PARAMETERS	INTENDED IMPROVEMENT
Shingles Vaccination Rate ≥50 YOA	% of adults aged 50 years and older who are vaccinated against shingles	<ul style="list-style-type: none"> <li>Person is ≥50 YOA as of today's date</li> </ul>	Increase the % of adults aged 50 years and older who have completed shingles vaccination*
Adult Immunization Status: AIS-E	% of eligible patients who are fully vaccinated against Zoster, Hepatitis B, Influenza, Pneumococcal and Td/Tdap	Measure for the Health Measure ID for AIS-E: Adult Immunization Status: Zoster* Measure Group Domain - Electronic Clinical Data Systems (ECDS) <b>NOTE:</b> Health systems should repeat this measure, replacing "Adult Immunization Status: Zoster" with each of the following: <ul style="list-style-type: none"> <li>Adult Immunization Status: Hepatitis B</li> <li>Adult Immunization Status: Influenza</li> <li>Adult Immunization Status: Pneumococcal</li> <li>Adult Immunization Status: Td/Tdap</li> </ul>	Improve overall AIS score as a result of increasing the % of patients immunized for shingles

\*2-dose series of SHINGRIX.

Table continues on next page →

Please see Important Safety Information for SHINGRIX on page 6 and full Prescribing Information, also available at [SHINGRIXHCP.com](http://SHINGRIXHCP.com).



## 09. How to Configure Registry Metrics/Key Performance Indicators to Monitor Population Health Metrics in Oracle Health (continued)

REGISTRY METRIC	KPI	RULE PARAMETERS	INTENDED IMPROVEMENT
Underperforming Sites	Sites that are below the shingles vaccination goal for this health system	<ul style="list-style-type: none"> <li>Rule to set the shingles vaccination goal AND Location (Facility, Unit, Building)</li> </ul> <p>(NOTE: the rule to set the shingles vaccination goal may already exist)</p>	Increase % of shingles vaccination for eligible patients across the underperforming site(s)
Shingles Vaccination Rate (IC patients ≥19 YOA and <50 YOA)	% of adults 19 through 49 YOA who are or will be at increased risk of HZ due to immunodeficiency or immunosuppression caused by known disease or therapy and who are vaccinated against shingles	<p><u>Option 1</u></p> <ul style="list-style-type: none"> <li>Person is ≥19 YOA as of today's date</li> </ul> <p>AND</p> <ul style="list-style-type: none"> <li>Person is &lt;50 YOA as of today's date</li> </ul> <p>AND</p> <ul style="list-style-type: none"> <li>Problem of IC State (not in a resolved or inactive status) at any time prior to the end of the current measurement period</li> </ul> <p>See pages 17-18 for examples of conditions to help identify IC patients: HIV/AIDS, Hematologic malignancy and Other immune conditions</p>	Increase the % of IC adults 19 through 49 YOA who have completed shingles vaccination*

\*2-dose series of SHINGRIX.

Table continues on next page →

Please see Important Safety Information for SHINGRIX on page 6 and full Prescribing Information, also available at [SHINGRIXHCP.com](http://SHINGRIXHCP.com).



## 09. How to Configure Registry Metrics/Key Performance Indicators to Monitor Population Health Metrics in Oracle Health (continued)

REGISTRY METRIC	KPI	RULE PARAMETERS	INTENDED IMPROVEMENT
Shingles Vaccination Rate (IC patients ≥19 YOA and <50 YOA)	% of adults 19 through 49 YOA who are or will be at increased risk of HZ due to immunodeficiency or immunosuppression caused by known disease or therapy and who are vaccinated against shingles	<p><u>Option 2</u></p> <ul style="list-style-type: none"> <li>Person is ≥19 YOA as of today's date</li> </ul> <p>AND</p> <ul style="list-style-type: none"> <li>Person is &lt;50 YOA as of today's date</li> </ul> <p>AND</p> <ul style="list-style-type: none"> <li>Problem of IC State (not in a resolved or inactive status) at any time prior to the end of the current measurement period</li> </ul> <p>See pages 18-19 for examples of conditions to help identify IC patients: Solid malignancy, Organ transplant and Rheumatologic/ inflammatory)</p> <p>AND</p> <p>Medication during the current measurement period</p> <p>See page 19 for examples of medications to help identify IC patients<sup>†</sup></p>	Increase the % of IC adults 19 through 49 YOA who have completed shingles vaccination*

\*2-dose series of SHINGRIX.

<sup>†</sup>For systemic corticosteroids, consider a parameter of >14 days of consecutive use.

Table continues on next page →

Please see Important Safety Information for SHINGRIX on page 6 and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](#).



## 09. How to Configure Registry Metrics/Key Performance Indicators to Monitor Population Health Metrics in Oracle Health (continued)

REGISTRY METRIC	KPI	RULE PARAMETERS	INTENDED IMPROVEMENT
Shingles Vaccination Rate (IC patients ≥19 YOA and <50 YOA)	% of adults 19 through 49 YOA who are or will be at increased risk of HZ due to immunodeficiency or immunosuppression caused by known disease or therapy and who are vaccinated against shingles	<p><u>Option 3</u></p> <p>Person is ≥19 YOA as of today's date</p> <p>AND</p> <p>Person is &lt;50 YOA as of today's date</p> <p>AND</p> <p>Medication during the current measurement period</p> <p>See page 19 for examples of medications to help identify IC patients<sup>†</sup></p>	Increase the % of IC adults 19 through 49 YOA who have completed shingles vaccination*
Shingles Vaccination Rate (≥50 YOA with certain chronic conditions)  NOTE: Health systems may run this metric for any single comorbidity or as a list based upon organizational preference	% of adults aged 50 years and older, with one or more certain comorbidities, who are vaccinated against shingles	<ul style="list-style-type: none"> <li>Person is ≥50 YOA as of today's date</li> </ul> <p>AND</p> <ul style="list-style-type: none"> <li>Problem of chronic condition(s) (not in a resolved or inactive status) at any time prior to the end of the current measurement period</li> </ul> <p>See pages 20-21 for information regarding comorbidity ICD-10 codes</p>	Increase the % of adults aged 50 years and older with one or more certain comorbidities who have completed shingles vaccination*

\*2-dose series of SHINGRIX.

<sup>†</sup>For systemic corticosteroids, consider a parameter of >14 days of consecutive use.

Please see Important Safety Information for SHINGRIX on page 6 and full Prescribing Information, also available at [SHINGRIXHCP.com](http://SHINGRIXHCP.com).



## 09. How to Configure Registry Metrics/Key Performance Indicators to Monitor Population Health Metrics in Oracle Health (continued)

Health system customers are encouraged to rely on existing Diagnosis Related Groupers (DRGs) within their EHR systems to identify specific patient populations.

For additional information, as needed, see below for a list of example diagnoses that may be associated with relevant comorbidities.

Health systems may use these example diagnoses to assist in potentially identifying the parameters for the search query.

**NOTE: The conditions and therapeutic agents listed in Tables 1 and 2 are not exhaustive and only serve as an example to help identify potentially immunocompromised patients. Health system stakeholders should review and amend as they see fit to identify and manage immunocompromised patients.**

**The degree of altered immunocompetence in a patient should be determined by a physician.<sup>2</sup>**

Please see the Background section of this document (Section 02) for an overview of the populations included in the clinical trials involving SHINGRIX.

Table 1. Conditions and ICD-10-CM Codes Used to Identify Enrollees With Immunosuppression in MarketScan Database Study, United States, July 2012–August 2017 <sup>3*</sup>	
Condition	ICD-10 codes
HIV/AIDS <sup>†</sup> HIV/AIDS disease	B20-B24
<b>Hematologic malignancy</b> Lymphatic and hematopoietic tissue malignancy	C81-C83; C88-C96
<b>Other immune conditions<sup>†</sup></b> Disorders of immune mechanism Neutropenia Functional disorders of neutrophils Genetic anomalies of leukocytes Decreased leukocyte count Leukocyte disease NEC <sup>§</sup> Leukocyte disease NOS Myelofibrosis Blood diseases NEC Blood diseases NOS Immunologic findings NEC Nonspecific immune findings NEC and NOS	D89 D70 D71 D72.0 D72.81 D72.89 D72.9 D75.81 D47.4; D75.89; D75.9; D89.2 D75.9; D75.89 R76; R83.4-R87.4; R89.4 R76; R83.4-R87.4; R89.4

Table continues on next page →

Please see Important Safety Information for SHINGRIX on page 6 and full Prescribing Information, also available at [SHINGRIXHCP.com](http://SHINGRIXHCP.com).



## 09. How to Configure Registry Metrics/Key Performance Indicators to Monitor Population Health Metrics in Oracle Health (continued)

**Table 1. Conditions and ICD-10-CM Codes Used to Identify Enrollees With Immunosuppression in MarketScan Database Study, United States, July 2012–August 2017<sup>3\*</sup>**

Condition	ICD-10 codes
<b>Solid malignancy</b> Organ/system malignant tumors Neuroendocrine tumors Neoplasms of uncertain behavior	C00-C07; C11-C19; C22-C80; Z85 C7A; C7B; D3A D00-D49
<b>Organ transplant<sup>†</sup></b> Complications of transplanted organ Organ transplant status	T86 Z94; Z98.85
<b>Rheumatologic/inflammatory<sup>#</sup></b> Sarcoidosis Amyloidosis NOS Familial Mediterranean fever Amyloidosis NEC Multiple sclerosis Other CNS demyelination Acute infective polyneuritis Acute myocarditis Polyarteritis nodosa and other Allergic alveolitis/pneumonitis NOS Other alveolar pneumonopathy Enteritis and colitis Lupus erythematosus Diffuse connective tissue disease Arthropathy with infection Crystal arthropathies Rheumatoid arthritis/inflammatory polyarthropathy Inflammatory spondylopathies Polymyalgia rheumatica	D86 E85 E85.0; M04 E85.1; E85.3; E85.8 G35 G36; G37.1; G37.3; G37.8; G37.9 G61.0; G61.9 I40 M30 T78.40; J67.9 J84.01; J84.02; J84.09 K50-K52 L93.0; L93.2; M32 L94; M35.8; M35.9 M12.9; M01.X0; M02.10 M11 M05-M14 M46 M31.5; M35.3

Table adapted from Patel M, Chen J, Kim S, et al. Analysis of MarketScan Data for Immunosuppressive Conditions and Hospitalizations for Acute Respiratory Illness, United States. *Emerg Infect Dis.* 2020;26(8):1720-1730.

\*Data from a national MarketScan Commercial Claims and Medicare database study from August 1, 2012, through July 31, 2017, estimating the prevalence of immunosuppressive conditions. The study used an algorithm to identify immunosuppressive conditions based on 6 groups of diseases and 3 classes of medications.

<sup>†</sup>Excludes asymptomatic HIV code of ICD-10 (Z21).

<sup>#</sup>Sickle cell disease, asplenia, and psoriatic arthritis were not included in the Greenberg algorithm but are considered to have immune deficiencies by Infectious Diseases Society of America guidelines. Adding these to the algorithm only increased the prevalence of immunosuppressive conditions by 0.1%.

<sup>§</sup>When a specific code is not available for a condition, the Index directs the coder to the “other specified” code in the Tabular List. NEC is used to indicate the diagnosis is specific; however, the coding system is not specific.

<sup>¶</sup>Bone marrow and peripheral stem cell transplant were considered under organ transplant and only considered immunosuppressed if enrollees were currently being given chemotherapeutic agents or immune modulators. Considering these enrollees under other immune conditions in which immunosuppressed does not require receipt of chemotherapeutic agents or immune modulators would increase the overall prevalence of immunosuppressed by 0.01%.

<sup>‡</sup>Psoriatic arthritis was not included in the Greenberg algorithm and could be an indication for immunosuppressive treatment. Adding this condition did not increase the prevalence of immunosuppressive conditions.

CNS, central nervous system; ICD-10-CM, International Classification of Diseases, 10th Revision, Clinical Modification; NEC, not elsewhere classified; NOS, not otherwise specified.

**Please see Important Safety Information for SHINGRIX on page 6 and full Prescribing Information, also available at [SHINGRIXHCP.com](http://SHINGRIXHCP.com).**



## 09. How to Configure Registry Metrics/Key Performance Indicators to Monitor Population Health Metrics in Oracle Health (continued)

Table 2. Examples of Medications Used to Identify Enrollees With Immunosuppression in a MarketScan Database Study, United States <sup>3*</sup>			
<b>Chemotherapeutic</b>	• Aldesleukin	• Dasatinib	• Pegaspargase
	• Alemtuzumab	• Decitabine	• Pemetrexed
	• Arsenic trioxide	• Denileukin diftitox	• Pentostatin
	• Azacitidine	• Docetaxel	• Pertuzumab
	• Bendamustine hydrochloride	• Etoposide	• Pralatrexate
	• Bevacizumab	• Everolimus	• Rituximab
	• Bexarotene	• Floxuridine	• Romidepsin
	• Bortezomib	• Fluorouracil	• Temozolomide
	• Brentuximab vedotin	• Gefitinib	• Thioguanine
	• Busulfan	• Ifosfamide	• Thiotepa
	• Cabazitaxel	• Ipilimumab	• Trastuzumab
	• Capecitabine	• Ixabepilone	• Tretinoin
	• Carboplatin	• Lomustine	• Vorinostat
	• Carfilzomib	• Melphalan	
	• Carmustine	• Mercaptopurine	<b>Immune-modulating</b>
	• Cetuximab	• Mesna	• Abatacept
	• Chlorambucil	• Methotrexate	• Adalimumab
	• Cisplatin	• Mitomycin	• Alefacept
	• Cladribine	• Mitotane	• Anakinra
	• Cyclophosphamide	• Nelarabine	• Auranofin
• Dacarbazine	• Ofatumumab	• Azathioprine	
• Dactinomycin	• Oxaliplatin	• Basiliximab	
	• Paclitaxel	• Belatacept	
	• Panitumumab	• Belimumab	
			• Certolizumab pegol
			• Cyclosporine
			• Daclizumab
			• Denosumab
			• Eculizumab
			• Etanercept
			• Golimumab
			• Infliximab
			• Interferon alfacon-1
			• Leflunomide
			• Lenalidomide
			• Mycophenolate mofetil
			• Natalizumab
			• Sirolimus
			• Tacrolimus
			• Thalidomide
			• Tocilizumab
			• Ustekinumab
			<b>Systemic corticosteroids<sup>†</sup></b>
			• Dexamethasone
			• Methylprednisolone
			• Prednisolone
			• Prednisone

**NOTE:** The degree to which immunosuppressive drugs cause clinically significant immunodeficiency generally is dose related and varies by drug. The degree of altered immunocompetence in a patient should be determined by a physician.<sup>2</sup>

Table adapted with changes from Patel M, Chen J, Kim S, et al. Analysis of MarketScan Data for Immunosuppressive Conditions and Hospitalizations for Acute Respiratory Illness, United States. *Emerg Infect Dis.* 2020;26(8):1720-1730.

\*Data from a national MarketScan Commercial Claims and Medicare database study from August 1, 2012, through July 31, 2017, estimating the prevalence of immunosuppressive conditions. The study used an algorithm to identify immunosuppressive conditions based on 6 groups of diseases and 3 classes of medications.

<sup>†</sup>Corticosteroids >14 days.

Please see Important Safety Information for SHINGRIX on page 6 and full Prescribing Information, also available at [SHINGRIXHCP.com](http://SHINGRIXHCP.com).



## 09. How to Configure Registry Metrics/Key Performance Indicators to Monitor Population Health Metrics in Oracle Health (continued)

Health system customers are encouraged to rely on existing DRGs within their EHR systems to identify specific patient populations.

The Wellness Registry will identify patients for SHINGRIX per ACIP recommendations. You may also want to educate these patients regarding their risk of shingles associated with certain chronic conditions. The following diagnosis codes may help identify those patients in your search query.

<b>Table 3. Examples of Comorbidities That Have Been Associated with an Increased Risk of HZ<sup>4</sup></b>	
<b>Disease</b>	<b>ICD-10 Code/Description</b>
COPD	J44.0 - Chronic obstructive pulmonary disease with (acute) lower respiratory infection J44.1 - Chronic obstructive pulmonary disease with (acute) exacerbation J44.9 - Chronic obstructive pulmonary disease, unspecified
Heart failure*	I50 - Heart failure
Heart disease*	I51.9 - Heart disease, unspecified
Hypertension*	I11 - Hypertensive heart disease I10 - Essential (primary) hypertension
Hyperlipidemia*	E78.0 - Pure hypercholesterolemia E78.1 - Pure hyperglyceridemia E78.2 - Mixed hyperlipidemia E78.3 - Hyperchylomicronemia E78.4 - Other hyperlipidemia E78.5 - Hyperlipidemia, unspecified

Table continues on next page →

Please see Important Safety Information for SHINGRIX on page 6 and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](#).



## 09. How to Configure Registry Metrics/Key Performance Indicators to Monitor Population Health Metrics in Oracle Health (continued)

<b>Table 3. Examples of Comorbidities That Have Been Associated with an Increased Risk of HZ<sup>4</sup></b>	
<b>Disease</b>	<b>ICD-10 Code/Description</b>
Stroke*	I63 - Cerebral infarction
Atrial fibrillation/flutter*	I48 - Atrial fibrillation and flutter
Chronic renal disease	N18 - Chronic kidney disease (CKD)
Asthma	J45 - Asthma
Diabetes	E08 - Diabetes mellitus due to an underlying condition E10 - Type 1 diabetes mellitus E11 - Type 2 diabetes mellitus

\*In a meta-analysis cardiovascular conditions were associated with an increased risk of HZ. Note, the cardiovascular conditions included in each individual study assessed in the meta-analysis varied and included heart disease, heart failure, hypertension, hyperlipidemia, stroke, atrial fibrillation/flutter, and other cardiovascular disease. These conditions when assessed individually may not be associated with an increased risk of HZ. Clinical judgment should be used when assessing risk factors for HZ.

Please see Important Safety Information for SHINGRIX on page 6 and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](#).



# 10. Key Population Health Configuration Use Cases

## 10.1 How to access and use the registry data/scorecards

Population health registry data can be accessed in the Oracle Health EHR in a few ways:

1. Reports via Dynamic Worklist or DA2
2. The HealthRegistries application

The below provides an overview and instructions for accessing each of the above-listed options.

### Reports

Dynamic Worklist or DA2 are end-user reporting tools that can be used to access registry data. Access and permission to either reporting tool may vary but both can be used to review patient information. A comprehensive patient search list and automated patient outreach resource provides full details on how to leverage both reporting solutions. All patients who meet the registry inclusion rule are included in the registries and can be included in the report. Consider the registry and the registry metric criteria to create the desired report. For example, to find all appropriate patients without a shingles immunization, use the Registry criterion and the Registry Metric count criterion for the number of shingles immunizations for the patient.

Registry-based information can be added in many reporting templates. Consider adding registry criteria (for example, the Registry Metric criterion) and display columns (for example, the Patient Registry Metric Value column) in reports. A unique Registry criterion is available in Dynamic Worklist.

Please see Important Safety Information for SHINGRIX on [page 6](#) and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](#).



## 10. Key Population Health Configuration Use Cases (continued)

### HealtheRegistries

1. **Access HealtheRegistries** in the app bar. Access may be restricted to certain roles.
2. **Click the Registries tab** on the top of the screen. **Consider selecting** Scorecard or Announcements to review summarized information if desired.
3. From the available options, **select an organization**. The provided options are determined by the initial configuration of the registry in HealtheIntent.
4. From the pulldown menu, **select the desired registry**, for example, “wellness”.
5. Registry Supporting Facts can be sorted by clicking the Completion % or Met % options in the pull-down menu. To view the data in a spreadsheet, **click the table button**. Registry Supporting Facts tiles that are red indicate that the measure is not achieved. The tile turns blue once the measure is achieved.
6. For individual provider metrics, **click the organization to reveal all associated providers**. Clicking on an individual provider reveals their performance against the selected Registry Fact.
7. **Click Export** to export the selected data set.

### 10.2 How to share population health reports and scorecards

Results from Dynamic Worklist and DA2 reports can be exported and shared to other users.

#### For Dynamic Worklist:

1. **Navigate to the List Actions tab** when creating a report.
2. **Select Share** and enter the user, group, or pool with whom to share the report.

#### For DA2:

1. **Navigate to the Report and right-click the name of the report**.
2. **Select Schedule from the drop-down menu** and select the user, group, or with whom to share the report.

Users can be provided access to dashboards within HealtheRegistries in order to access the application.

Please see **Important Safety Information for SHINGRIX on page 6** and full **Prescribing Information**, also available at [SHINGRIXHCP.com](http://SHINGRIXHCP.com).



# 10. Key Population Health Configuration Use Cases (continued)

## 10.3 How to create immunization tactics in Oracle Health's HealthIntent

Because Population Health touches many stakeholders in a health system, gaining buy-in from the key constituents is critical. Building a shingles immunization program requires a strategy and approved tactics.

When setting up the shingles population health program, consider the following milestones:

- 1.** Identify all stakeholders: The population health/quality team may have extensive knowledge in setting up similar programs and have a playbook available. A clinical champion/leadership team tasked to drive immunizations across the health system can orchestrate the strategy and tactics. The EHR team may need to refine the registry inclusion rule and registry metrics and create reports and clinical decision support aligned to the strategy.
- 2.** Creating the shingles immunization program: A timeline of milestones, objectives, and key personnel involved can set expectations for each phase of the shingles program.
- 3.** An interface may be required to connect with the state immunization registry to provide a complete immunization history of patients.
- 4.** Identify the target population: Given historical shingles immunization rates, what is the target rate to achieve?
- 5.** Besides the overall shingles immunization rate, what other key performance (registry) metrics related to shingles can be defined and are meaningful?

Please see Important Safety Information for SHINGRIX on [page 6](#) and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](#).



## 11. Disclaimers

- Conditions, ICD codes, and medications listed in this guide are only examples to help identify patients. Ultimately it is up to the HCP/health system to select the conditions, codes, and/or medications they feel put a patient at increased risk of HZ and to determine whether a patient is appropriate for SHINGRIX
- The conditions and therapeutic agents listed in Tables 1 and 2 are not exhaustive and only serve as an example to help identify potentially immunocompromised patients. Health system stakeholders should review and amend as they see fit to identify and manage immunocompromised patients. The degree of altered immunocompetence in a patient should be determined by a physician
- The customer (ie, physician, medical group, IDN) shall be solely responsible for the implementation, testing, and monitoring of the instructions to ensure proper orientation in each customer's EHR system
- Capabilities, functionality, and setup (customization) for each individual EHR system vary. GSK shall not be responsible for revising the implementation instructions it provides to any customer if the customer modifies or changes its software or the configuration of its EHR system, after such time as the implementation instructions have been initially provided by GSK
- While GSK tests its implementation instructions on multiple EHR systems, the instructions are not guaranteed to work for all available EHR systems and GSK shall have no liability thereto
- While EHRs may assist providers in identifying appropriate patients for consideration of assessment, treatment, and referral, the decision and action should ultimately be decided by a provider in consultation with the patient, after a review of the patient's records to determine eligibility, and GSK shall have no liability thereto
- The instructions have not been designed to and are not tools or solutions for meeting Advancing Care Information and/or any other quality/accreditation requirement
- All products are trademarks of their respective holders, all rights reserved. Reference to these products is not intended to imply affiliation with or sponsorship of GSK and/or its affiliates

**Please see Important Safety Information for SHINGRIX on [page 6](#) and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](#).**





Please see Important Safety Information for SHINGRIX on page 6 and full [Prescribing Information](#), also available at [SHINGRIXHCP.com](#).

**References:** **1.** Prescribing Information for SHINGRIX. **2.** General Best Practices for Immunization. Centers for Disease Control and Prevention. July 25, 2024. Accessed June 13, 2025. <https://www.cdc.gov/vaccines/hcp/imz-best-practices/index.html> **3.** Patel M, Chen J, Kim S, et al. Analysis of MarketScan data for immunosuppressive conditions and hospitalizations for acute respiratory illness, United States. *Emerg Infect Dis.* 2020;26(8):1720-1730. **4.** Marra F, Parhar K, Huang B, Vadlamudi N. Risk factors for herpes zoster infection: a meta-analysis. *Open Forum Infect Dis.* 2020;7(1):1-8.

Trademarks are property of their respective owners.

©2025 GSK or licensor.

PMUS-SGXLBND250039 December 2025

Produced in USA.