

STARTING AT AGE 50, YOUR RISK OF SHINGLES SHARPLY RISES

You are at increased risk of shingles due to (check all that apply):

√ 50+ YEARS	ASTHMA	COPD CARDIOVASCULAR CONDITIONS
DIABETES	CHRONIC KIDNEY DISEASE	OTHER
		(Other reason determined by your pharmacist)

COPD=chronic obstructive pulmonary disease.

UNDERSTANDING THE IMPORTANCE OF SHINGLES PREVENTION

What is shingles?

Shingles is a **painful, blistering rash** that can last for weeks and disrupt work, sleep, and time with family.

Shingles is caused by the same virus that causes chickenpox. The virus stays inside your body after you have chickenpox and can reactivate at any time, causing shingles. **More than 99% of people 50 years and older** have the virus that causes shingles inside them. One in 3 people will develop

What is SHINGRIX?

shingles in their lifetime.

SHINGRIX is an **FDA-approved vaccine** for the prevention of shingles (herpes zoster) in adults 50 years and older. SHINGRIX is not used to prevent chickenpox.

What should I know about SHINGRIX?

SHINGRIX demonstrated outstanding efficacy. In clinical trials, SHINGRIX was proven to be **more than 90% effective** in preventing shingles in adults 50 years and older who completed the 2-dose series.

Get the **full protection** offered by SHINGRIX by completing the 2-dose series.

The **most common side effects of SHINGRIX** are pain, redness, and swelling at the injection site, muscle pain, tiredness, headache, shivering, fever, and upset stomach.

What does SHINGRIX cost?

\$0 for almost everyone: \$0with Medicare Part D and \$0 for 98% of privately insured people.







Learn more about shingles at SHINGRIX.com

†Coverage and cost may vary and are subject to change without notice. Reimbursement decisions are made by individual insurance plans.

Important Safety Information

 You should not receive SHINGRIX if you are allergic to any of its ingredients or had an allergic reaction to a previous dose of SHINGRIX

Please see next page for additional Important Safety Information about SHINGRIX.





^{*}Cardiovascular conditions include heart disease, heart failure, hypertension, hyperlipidemia, stroke, atrial fibrillation/flutter, and other cardiovascular disease.

SHINGLES CAN DISRUPT YOUR ROUTINE **HELP PROTECT YOURSELF AGAINST** SHINGLES WITH SHINGRIX

- In clinical trials, SHINGRIX was proven to be more than **90% effective in preventing shingles** in adults 50 years and older who completed the 2-dose series.
- Pain, redness, and swelling at the injection site, muscle pain, tiredness, headache, shivering, fever, and upset stomach are the most common side effects. In studies, these side effects had a median duration of 2 to 3 days.
- You can get SHINGRIX at the same time you get certain other vaccines, such as your flu shot. Talk to your doctor or pharmacist.
- SHINGRIX is given as a 2-dose series. To get the full protection offered by SHINGRIX, it's important to receive both doses. After your first dose, the second dose should be administered 2 to 6 months later.



Scan the QR code or visit ScheduleSHINGRIX.com to book both doses of your SHINGRIX vaccination. You can also get a reminder for your second dose at SHINGRIX reminder.com

Track your appointments by writing them below:

Important Safety Information (cont'd)

- An increased risk of Guillain-Barré syndrome (severe
 SHINGRIX was not studied in pregnant or nursing muscle weakness) was observed after vaccination with SHINGRIX
- Fainting can happen after getting injectable vaccines, including SHINGRIX. Precautions should be taken to avoid falling and injury due to fainting
- The most common side effects are pain, redness, and swelling at the injection site, muscle pain, tiredness, headache, shivering, fever, and upset stomach
- women. Tell your healthcare provider if you are pregnant, plan to become pregnant, or are breastfeeding
- Vaccination with SHINGRIX may not protect all individuals
- Ask your healthcare provider about the risks and benefits of SHINGRIX. Only a healthcare provider can decide if SHINGRIX is right for you

You are encouraged to report vaccine adverse events to the US Department of Health and Human Services. Visit www.vaers.hhs.gov to file a report or call 1-800-822-7967. You may also report negative side effects to GSK at https://gsk.public.reportum.com or 1-888-825-5249.

Please see additional Important Safety Information on the previous page and **Important Facts About SHINGRIX.**

